



Introduction to Population Grids and their Integration with Remote Sensing Data for Sustainable Development and Disaster Management

April 6, 2021

# NASA's Applied Remote Sensing Training Program (ARSET)

https://appliedsciences.nasa.gov/what-we-do/capacity-building/arset/about-arset

- Part of NASA's Applied Sciences Program
- Empowering the global community through remote sensing training
- Seeks to increase the use of Earth science in decision-making through training for:
  - Policy makers
  - Environmental managers
  - Other professionals in the public and private sector





#### The POPGRID Data Collaborative

https://www.popgrid.org/about-us





BILL& MELINDA GATES foundation







Center for International Earth Science Information Network EARTH INSTITUTE | COLUMBIA UNIVERSITY



## **Course Structure and Materials**

- Two, 2-hour sessions on March 30 and April 6
- The same content will be presented at two different times each day:
  - Session A: 10:00-12:00 EDT (UTC-4)
  - Session B: 15:00-17:00 EDT (UTC-4)
  - You only need to attend one session per day.
- Webinar recordings, PowerPoint presentations, and the homework assignment can be found at:
  - <u>https://appliedsciences.nasa.gov/join-</u> <u>mission/training/english/arset-introduction-</u> <u>population-grids-and-their-integration-remote</u>
  - Q&A following each lecture and/or by email at:
    - <u>brock.blevins@nasa.gov</u>
    - <u>nasa.arset@gmail.com</u>



HOME / JOIN THE MISSION / TRAINING



## **Homework and Certificates**

- Homework:
  - One homework assignment given after Part 2
  - Answers must be submitted via Google
     Forms
  - HW Deadline: Tuesday April 27


#### • Certificate of Completion:


- Attend both live webinars (March 30 and April 6)
- Complete the homework assignment by the deadline (access from ARSET website)
- You will receive certificates approximately three months after the completion of the course from: <u>marines.martins@ssaihq.com</u>






#### **Course Outline**











# Introduction to the Applications of Population Grids

#### Dr. Robert S. Chen


Director and Senior Research Scientist, CIESIN Manager, NASA Socioeconomic Data and Applications Center Co-chair, Thematic Research Network on Data and Statistics (TReNDS) UN Sustainable Development Solutions Network

April 06, 2021




# **Original Gridded Population of the World Dataset**

- Created in 1994-95
  - Collaboration with NCGIA (Waldo Tobler and Uwe Deichmann)
  - 19,000 administrative units used
  - Estimated 1994 populations; total population of 5.6 billion
  - 5' x 5' lat-lon spherical quadrilaterals, 6.7 million cells
  - Unsmoothed and smoothed versions (Tobler's pycnophylactic algorithm)
  - Regional and global coverages






#### Gridded Population of the World – Version 2




## GPWv4.11, Population by Age Group & Gender, 2010



GPWv4.11 includes a dataset on Basic Demographic Characteristics with gridded estimates of population by age groups and gender for 2010.


#### http://sedac.ciesin.columbia.edu/data/coll ection/gpw-v4/sets/browse



Gridded Population of the World, Version 4 (GPWv4) Basic Demographic Characteristics, Revision 10 consists of estimates of human population by age and sex, consistent with national censuses and population registers, for the year 2010. The sex ratio represents the number of males for every 100 females in the population. It is calculated by dividing the total male count rater by the total female count rater and multiphying the quotient by 100.

Basic Demographic Characteristics, v4.10, 2010: Women of Childbearing Age (Ages 15-49)


Gridded Population of the World, Version 4 (GPWv4)





Gridded Population of the World, Version 4 (GPWW4) Basic Demographic Characteristics, Revision 10 consists of estimates of human population by age and sex, consists national censuses and population registers, for the year 2010. The global distribution of women of childbearing age, ages 15–69, is represented here as a percent of the population in the year 2010. It is calculated by summing the counts in the v4.10 female 5-year age group rasters from ages 15 to 40, then dividing the sum by the total popul the year 2010 and mulbiplying the quotient by 100.


# New Sources of Data for Mapping Human Settlements, Infrastructure, and Population

- Night-Time Lights (DMSP >1 km  $\rightarrow$  VIIRS ~750 m)
- Landsat (~30 m)
- Radar (~12 m)
- High Resolution Imagery (< ~3 m resolution)</li>











Center for International Earth Science Information Network EARTH INSTITUTE | COLUMBIA UNIVERSITY

## High Resolution Settlement Layer from Facebook

- Result of a collaboration between Facebook, CIESIN, and the World Bank
- Originally motivated by the desire to optimize Internet access in rural areas
- Further development driven by humanitarian needs identified by the IFRC and others



ganization/facebook



https://tech.fb.com/ai-powered-maps-helpvaccination-campaigns/

NASA's Applied Remote Sensing Training Program

#### **POPGRID** Website and Viewer








How the POPGRID Data Community is Responding to COVID-19 - 4 June 2020 Gap As communities around the world grapple with the devastating effects

A New POPGRID Report Cities, crowding, and the Shows How Gridded Popcoronavirus: Predicting ulation Data Can Help contagion risk hotspots Close the Knowledge Data - 12 May 2020 This blog post is reposted from the

World Bank's Sustainable Cities blog

https://www.popgrid.org/

- 3 June 2020



#### https://sedac.ciesin.columbia.edu/map ping/popgrid/

Center for International Earth Science Information Network EARTH INSTITUTE | COLUMBIA UNIVERSITY

## Applications of SEDAC's Gridded Population Data in Scientific Literature


- More than 3,300 citations of GPW versions in SEDAC's citation database
- More than one-fourth of these citations also cite remote sensing data

	omic Data ani		IS CENTER (SEDAC) - Hosted by <u>CIESIN</u> at Columbia University	
		4	ويوالد انتهر فواليس	
ATA MAPS	THEMES	RESOURCES	Social Media ABOUT	
	5 THEMES	RESOURCES .	SOCIAL MEDIA	
Citations Database			Follow Us: 🐸 🖪 💶 💶   Shi	
e Citations Database is a collection	of all identified publications the	at cite SEDAC data.		
Filter by:	All Fields		0	
facets are ordered by results count)			<u> </u>	
Uses Remote Sensing?	Title	Abstract	First author	
Yes (948)				
	Sort by Order	Items per page	y Reset	
Year	Year • Descendi	ng • 10 •	( Treaser	
2020 (427)	Displaying 1 - 10 of 3377			
2018 (414)	Export results: CSV XML	OR Ex	port selected citations using checkboxes: CSV/XML	
2019 (388)	Abel, Christin: Horion, Sté	phanie: Tagesson, Torbern: D	e Keersmaecker, Wanda; Seddon, Alistair W. R.;	
2017 (347)			man-environment nexus and vegetation-	
2016 (321)	rainfall sensitivity in trop	oical drylands. [Nature Susta	inability] . 4(25-32) DOI:	
Show more	https://doi.org/10.1038/s41	893-020-00597-z		
Publication type	Uses Remote Sensing: yes			
<ul> <li>Journal Article (2765)</li> </ul>	SEDAC Data Collection(s): • g	pw-v4 • wildareas-v3	(Journal Article) export	
<ul> <li>Report (257)</li> </ul>	Bai, Heming; Zheng, Zhi;	Zhang, Yuanpeng; Huang, He	; Wang, Li. 2021. Comparison of satellite-based	
<ul> <li>Book Section (190)</li> </ul>	PM2.5 estimation from a	erosol optical depth and top	-of-atmosphere reflectance. [Aerosol and Air	
Conference Proceedings		200257 DOI: https://doi.org/1	0.4209/aaqr.2020.05.0257	
(33)	Uses Remote Sensing: yes			
<ul> <li>Book (29)</li> </ul>	SEDAC Data Collection(s): • g	pw-v4	(Journal Article) export	
Show more	Bayat, Bagher; Camacho, I	Fernando; Nickeson, Jaime; C	Cosh, Michael; Bolten, John; Vereecken, Harry;	
SEDAC Data Collections	Montzka, Carsten. 2021. T	oward operational validatio	n systems for global satellite-based terrestrial oplied Earth Observation and Geoinformation]. 95:	
✓ gpw-v3		g/10.1016/j.jag.2020.102240	pare bara observation and ocomornation]. 35.	
✓ grump-v1	Uses Remote Sensing: yes			
grump +1	SEDAC Data Collection(s): • g	rump-v1	(Journal Article) export	

#### https://sedac.ciesin.columbia.edu/citations-db

Center for International Earth Science Information Network Earth Institute | Columbia University

# 63 Subject Areas of Journals with Articles Co-Citing SEDAC Data with Remote Sensing Data, 2007-2016



Agriculture

- Area Studies
- Astronomy & Astrophysics
- Biochemistry & Molecular Biology
- Biodiversity & Conservation
- Biotechnology & Applied Microbiology
- Business & Economics
- Cell Biology
- Computer Science
- Demography
- Education & Educational Research
- Emergency Medicine
- Endocrinology & Metabolism
- Energy & Fuels
- Engineering
- Entomology
- Environmental Sciences & Ecology
- Evolutionary Biology

Downs, Chen, & Schumacher, AGU, 2017

#### **Recent Citations: Science**

- Lau, W. W. Y., Shiran, Y., Bailey, R. M., Cook, E., Stuchtey, M. R., Koskella, J., . . . Palardy, J. E. Evaluating scenarios toward zero plastic pollution. Science, First release, 23 July.
  - -Uses **GPW v4.11** in combination with HydroSHEDS to assess population within 1 km of water and to delineate urban from rural areas (using European Commission definition). HydroSHEDs is based on SRTM data.


		living in proximity vater	Proportion of population living in proximity to water				
Archetype	< 1 km	> 1 km	< 1 km	> 1 km			
HI-U	387,364,176	502,326,123	43.5%	56.5%			
HI-R	139,551,400	200,098,923	41.1%	58.9%			
UMI-U	738,055,814	953,720,947	43.6%	56.4%			
UMI-R	333,681,550	481,968,707	40.9%	59.1%			
LMI-U	957,825,466	1,115,472,519	46.2%	53.8%			
LMI-R	340,003,878	470,101,017	42.0%	58.0%			
LI-U	91,908,375	136,868,144	40.2%	59.8%			
LI-R	159,418,874	278,735,508	36.4%	63.6%			

https://doi.org/10.1126/science.aba9475



Impact Factor (2019): 41.845

- Lecocq, T., Hicks, S. P., Van Noten, K., van Wijk, K., Koelemeijer, P., De Plaen, R. S. M., . . . Xiao, H. Global quieting of high-frequency seismic noise due to COVID-19 pandemic lockdown measures. Science, First release, 23 July.
  - -Uses **GPW v4.11** to distinguish seismic measurements in dense vs. sparse areas.



https://doi.org/10.1126/science.abd2438

Fig. 1 Worldwide seismic station locations. Locations of the 268 global seismic stations with usable data (e.g., no long data gaps, working sensors) we analyzed. Lockdown effects are observed (red) at 185 out of 268 stations. Symbol size is scaled by the inverse of population density (<u>28</u>) to emphasize stations located in remote areas...

> Center for International Earth Science Information Network EARTH INSTITUTE | COLUMBIA UNIVERSITY

## **Recent Citations: Nature Communications**

- Boulange, J., Hanasaki, N., Yamazaki, D., & Pokhrel, Y. (2021). Role of dams in reducing global flood exposure under climate change. Nature Communications, 12(1), 417.
- Uses GPWv4.11 for population exposure estimates
- Also uses Global Reservoirs and Dams (GRaND) database (but not necessarily SEDAC's version)




Fig. 1: Population exposure to the historical 100-year river flood (keeping population constant at 2010 level) downstream of dams

https://doi.org/10.1038/s41467-020-20704-0

NASA's Applied Remote Sensing Training Program

# communications Impact Factor (2019): 12.121

#### **Recent Citations: The Lancet**

- Watts et al. 2020. The 2020 report of The Lancet Countdown on health and climate change: responding to converging crises. Online First, 2 December.
  - -Uses **GPWv4** in computing multiple health impact indicators including excess heat, wildfire risk, labor force impacts, and urban green space




Figure 1. Change in days of heatwave exposure relative to the 1986–2005 baseline in people older than 65 years

https://doi.org/10.1016/S0140-6736(20)32290-X




Figure 4. Population-weighted average changes in the number of days of exposure to very high or extremely high risk of wildfire in 2016–19 compared with 2001–04





Figure 40: Urban Green Space by WHO Region.

Center for International Earth Science Information Network EARTH INSTITUTE | COLUMBIA UNIVERSITY

## **Recent Citations: The Lancet**

- Sartorius, B., Cano, J., Simpson, H., Tusting, L. S., Marczak, L. B., Miller-Petrie, M. K., . . . Pullan, R. L. (2021). Prevalence and intensity of soil-transmitted helminth infections of children in Sub-Saharan Africa, 2000–18: a geospatial analysis. The Lancet Global Health, 9(1), e52-e60.
  - -Uses GPWv4.11 estimates for children aged 5-14 years to estimate population counts of infection and moderate-to-heavy infection by species


https://doi.org/10.1016/S2214-109X(20)30398-3




THE LANCET Impact Factor (2019): 60.392

Figure 1. Prevalence of any soil-transmitted helminth infections (A) and of moderate-to-high intensity infections (B), and the absolute number of prevalent cases (C), in children aged 5–14 years in sub-Saharan Africa, 2000–18

#### SEDAC Data Reuse: Global Forest Watch and PREPdata



The Global Human Settlement Layer population (GHS-POP) data developed with the JRC, and Global Reservoir and Dam (GRaND) data are available as layers in the Global Forest Watch interactive map, in conjunction with other satellite data and imagery <u>https://www.globalforestwatch.org/</u> NASA's Applied Remote Sensing Training Program



- Partnership for Resilience and Preparedness (PREP)
  - Map-based, open data online platform for climate adaptation and resilience planning
  - Includes >4 data sets from SEDAC, including GPWv4, GRanDv1, GHSL-POP, and Global Roads Open Access Dataset (gROADS)

#### https://www.prepdata.org/explore



# **Use of the SEDAC Population Estimation Service in the GeoCollaborate Decision Support Tool**

Streets \$

20190827951PRda

Florida20190828

NWS Wind Potential Fcst-ORL


Wind damage potential guidance for land-falling tropical systems

Layers

- Use open services to enable data and queries to be accessed via existing decision support systems
- Support flexible platforms that integrate and transform data from multiple sources into timely and usable information



NASA's Applied Remote Sensing Training Program




Science Information Network

Earth Institute | Columbia University

# **SEDAC Data Reuse: Informal Education**

- Several SEDAC and NASA datasets featured prominently in "Countryside" Exhibit at the Guggenheim Museum through February 15, 2021
  - GPW, Human Footprint/Last of the Wild, gROADS, Nuclear Power Plants
  - Nighttime lights, land surface temp, etc.





Center for International Earth Science Information Network EARTH INSTITUTE | COLUMBIA UNIVERSITY

# GEO Focused on the Role of Earth Observations to Address the Sustainable Development Goals

 "Population distribution" and "Cities and Infrastructure Mapping" are important to indicators and decision making related to all 17 goals.

> Alignments of the Goals with Specific Types of Earth Observations and Geospatial Information





SUSTAINABLE DEVELOPMENT GOALS		Population distribution	Cities and infrastructure mapping	Elevation or Loopraphy	Land cover and use mapping	Oceanographic observations	Hydrological and water quality observations	Atmospheric and air quality monitoring	Biodiversity and ecosystem observations	Agricultural monitoring	Hazards, disasters and environmental impact monitoring
1 No poverty											
2 Zero hunger											
3 Good health and well-being											
4 Quality education											
5 Gender equality											
6 Clean water and sanitation											
7 Affordable and clean energy											
8 Decent work and economic grow	h										
9 Industry, innovation and infrastructure											
10 Reduced inequalities											
11 Sustainable cities and communitie											
12 Responsible consumption and production											
13 Climate action											
14 Life below water											
15 Life on land	1										
16 Peace, justice and strong institutions											
17 Partnerships for the goals							Ce	nter	for	Inte	ernat
							Sci	ence	: Inf	orm	atio

onal Earth Network

EARTH INSTITUTE | COLUMBIA UNIVERSITY

# Role of Fundamental Geospatial Data Layers in Assessing SDG Indicators



No.	Indicator
1.5.1	Number of deaths, missing persons and persons affected by disaster per 100,000 people
7.1.1	Proportion of population with access to electricity
9.1.1	Proportion of the rural population who live within 2 km of an all-season road
11.1.1	Proportion of urban population living in slums, informal settlements or inadequate housing
11.2.1	Proportion of population that has convenient access to public transport
11.3.1	Ratio of land consumption rate to population growth rate
11.6.2	Annual mean levels of fine particulate matter (e.g., PM2.5 and PM10) in cities (pop. weighted)
11.7.1	Average share of the built-up area of cities that is open space for public use for all, by sex, age and persons with disabilities




Many SDG indicators directly require fundamental population, settlement, and infrastructure data.

- Consistency in methods between data is therefore important.
- Choice of datasets has implications for indicator accuracy and consistency.

# Earth Observations Toolkit for Sustainable Cities and Human Settlements


- Collaboration between UN-Habitat and GEO EO4SDGs
- More than 12 SEDAC datasets and 3 SEDAC interactive tools are featured
- Aimed at urban and national planners and statisticians
- Launch event on January 21, 2021 at UNSD side event









#### https://eo-toolkit-guo-un-habitat.opendata.arcgis.com/













Measuring Development Progress: Proportion of the Rural Population who Live within 2 km of an All-Season Road (SDG 9.1.1) Linda Pistolesi and James Gibson, CIESIN

Linda Fisiolesi ana James Gibson, C

April 6, 2021

### **Learning Objectives**

27

By the end of this session, you will be able to...

- 1. Describe Indicator 9.1.1 and how it is measured
- 2. Identify the geospatial inputs used to calculate the Rural Access Index (RAI)
- 3. Explain why the choice of gridded population input can lead to different results for this "simple" indicator
- 4. Recognize the advantages and the limitations of using currently available open global data sources to calculate RAI



## Sustainable Development Goal (SDG) 9

27

Build resilient infrastructure, promote inclusive and sustainable industrialization, and foster innovation.

**Target 9.1:** ...development of a quality, reliable, and resilient transport network....


#### Indicator 9.1.1 - The Rural Access Index (RAI)

Proportion of the rural population who live within 2 km of an all-season road



The Rural Access Index (RAI), World Bank Version 1

- Rural Access Index: A Key Development Indicator
   World Bank Transport Unit technical paper by Roberts et al. (2006)
- Approaches to measuring RAI:
  - Preferred: Household surveys with questions about access to transport
  - Alternative: GIS methods and model estimations where data inputs were available
- RAI Computed for 173 countries at the national level only.








The Rural Access Index (RAI), World Bank Version 2


#### Measuring Rural Access: Using New Technologies, 2016 Transport & ICT Report

#### Figure AI.1: Basic Methodological Framework






World Bank Geospatial RAI Data Inputs and Sources



The Rural Access Index (RAI), World Bank Version 2

Advantages of the Geospatial Methodology:

- More cost-effective and sustainable; no need to wait for the next survey
- More consistent across countries
- Allows for subnational RAI estimation - aiding prioritization of rural road improvements and monitoring
- 25 countries completed thus far



RAI (%) based on World Bank v2

🗖 0 - 20 📕 20 - 40 📕 40 - 60 🧾 60 - 80 🦲 80 - 100 📰 RAI Not Computed


# A Global Open Data Approach to the RAI

Advantages of using Open Global Datasets to Calculate RAI for Every Country:


- **Common inputs** enable better cross-country comparison
  - Common gridded population estimates several choices
  - Standardized administrative boundaries
  - Common urban/rural definition based on population density, counts, & satellite-derived built-up areas
  - Common road classification; continuously updated
- Automated script for faster processing and ease of repeatability
- Temporal comparison possible as datasets are updated



## A Global Open Data Approach to the RAI





# Population Input: Three Gridded Population Datasets



🔗 Sync Panels 💟 | 🎛 Repeat Map | 🚱 Zoom Out | 🗐 Legend | 🚯 Overview | 👯 Comparison View

- Global Human Settlement Population 2015 (GHS-POP)
- WorldPop 2019 (WP)
- High Resolution Settlement Layer 2015 (HRSL)






<u>Popgrid Viewer</u>: https://sedac.ciesin.columbia.edu/mapping/popgrid/

## **Population Inputs Compared**

Dataset	Sourc e	Concept/Method	Resolution
<u>Global Human</u> <u>Settlement</u> <u>Population 2015</u> (GHS-POP)	JRC & CIESIN	<ul> <li>Lightly modeled</li> <li>Binary dasymetric: Proportional allocation of census population to Landsat-derived built up areas</li> </ul>	1 km (30 arc-seconds)
<u>WorldPop</u> <u>Unconstrained</u> 2019 UN Adjusted (WP)	WorldPop	<ul> <li>Highly modeled</li> <li>Random Forest model and dasymetric redistribution: Census population allocated to grid cells based on statistical weighting of co-variate layers, e.g., urban extents, land cover, roads</li> </ul>	1km (30 arc-seconds)
High Resolution Settlement Layer 2015 (HRSL) NASA's Applied Remote Ser	Facebook & CIESIN	<ul> <li>Lightly modeled</li> <li>Binary dasymetric: Proportional allocation of census population to satellite-derived "settlement extents" defined by the presence of buildings</li> <li>m Source: https://www.popg</li> </ul>	30 m (1 arc-second) grid.org/explore-data




# Admin Input: Database of Global Administrative Areas (GADM)





# Rural Areas Input: GHS Settlement Model (GHS-SMOD)



https://ghsl.jrc.ec.europa.eu/ghs\_smod2019.php



#### Roads Input: OpenStreetMap (OSM)





Global OSM roads data was downloaded as a geopackage from: <u>https://download.osmdata.xyz</u>

#### Global openstreetmap data extracts | osmdata.xyz

Introduction: This project provides global data extracts based on OpenStreetMap data as GeoPackages. Each extract represents its related primary feature regarding the OpenStreetMap project.

Information about this project and data processing: osmdata.xyz on GitHub

Updates are available every month. Building is updated every odd month.

#### **Downloads**

Note: The GeoPackages are compressed into ZIP-Archives. Projection of the geodata: WGS 84 | EPSG 4326

Name	File	ZIP size MB approx.	GeoPackage size MB approx.	Last update dd.mm.yyyy
Address	address_EPSG4326.zip	13.682	49.009	04.01.2021
Aerialway	aerialway_EPSG4326.zip	7	32	04.01.2021
Aeroway	aeroway_EPSG4326.zip	82	253	04.01.2021
Amenity	amenity_EPSG4326.zip	2.665	8.547	04.01.2021



## Roads Input: OpenStreetMap (OSM)

#### **OpenStreetMap Data Extracts**

The OpenStreetMap data files provided on this server do **not** contain the user names, user IDs and changeset IDs of the OSM objects because these fields are assumed to contain personal information about the OpenStreetMap contributors and are therefore subject to data protection regulations in the European Union. Extracts with full metadata are available to OpenStreetMap contributors only.


Welcome to Geofabrik's free download server. This server has data extracts from the <u>OpenStreetMap project</u> which are normally updated every day. Select your continent and then your country of interest from the list below. (If you have been directed to this page from elsewhere and are not familiar with OpenStreetMap, we highly recommend that you read up on OSM before you use the data.) This open data download service is offered free of charge by Geofabrik GmbH.

Willkommen auf dem Geofabrik-Downloadserver. Hier gibt es Daten-Auszüge aus dem <u>OpenStreetMap-Projekt</u>, die normalerweise täglich aktualisiert werden. Wählen Sie aus dem Verzeichnis unten den Kontient und ggf. das Land, für die Sie Daten benötigen. (Wenn Sie von anderswo auf dieser Seite gelandet sind und von OpenStreetMap nichts wissen, dann ist es empfehlenswert, sich mit dem Projekt vertraut zu machen, bevor Sie mit den Daten arbeiten.) Diese Downloads werden von der Geofabrik GmbH kostenlos angeboten.

Click on the region name to see the overview page for that region, or select one of the file extension links for quick access.

Sub Region	Quick Links			
	.osm	n.pbf	.shp.zip	.osm.bz2
Africa	[.osm.pbf]	(4.3 GB)	×	[.osm.bz2]
Antarctica	[.osm.pbf]	(29.1 MB)	[.shp.zip]	[.osm.bz2]
Asia	[.osm.pbf]	(9.2 GB)	×	[.osm.bz2]
Australia and Oceania	[.osm.pbf]	(845 MB)	×	[.osm.bz2]
Central America	[.osm.pbf]	(432 MB)	×	[.osm.bz2]
Europe	[.osm.pbf]	(22.9 GB)	×	[.osm.bz2]
North America	[.osm.pbf]	(10.1 GB)	×	[.osm.bz2]
South America	[.osm.pbf]	(2.3 GB)	×	[.osm.bz2]

Technical details about this download service.



Solution in the second second

Nicht das Richtige dabei? Die Geofabrik ist ein auf OpenStreetMap spezialisiertes Beratungs- und Softwareentwicklungsunternehmen in Karlsruhe. Gern helfen wir Ihnen bei der Datenaufbereitung, Datenkonvertierung, Serverinstallation und ähnlichen Aufgaben. <u>Besuchen Sie unsere Webseite</u> und sprechen Sie mit uns, wenn wir Ihnen helfen können. Alternatively, country level OSM roads data can be downloaded from Geofabrik Download Server: <u>https://download.geofabrik.de</u>



41

# Roads Input: OpenStreetMap (OSM)





<b>Osmdata.xyz</b> https://github.com/michaelmgis/osmdata.xyz	<b>Geofabrik</b> https://download.geofabrik.de/
- Global extract	- Country extract
<ul> <li>Primary feature (e.g., highways) and all attributes</li> </ul>	<ul> <li>Selection of features (e.g., land use, roads, waterways) and attributes</li> </ul>
- Geopackage	- Shapefile
- Universal timestamp for the	<ul> <li>Different timestamps for each country</li> </ul>




# Defining OSM "All-Season Roads"


#### Two Definitions Based on OSM "fclass" Attribute

OSM fclass	World Bank	CIESIN
Primary & Primary Link	✓	~
Secondary & Secondary Link	✓	~
Tertiary & Tertiary Link	✓	~
Trunk	✓	~
Motorway	✓	~
Unclassified	-	~
Track Grade 1	-	~
Unknown	-	~
Road	-	~


#### World Bank Definition




#### **CIESIN** Definition



### **Global Open Data RAI Script**



# Global Open Data RAI Script (Continued)



NASA's Applied Remote Sensing Training Program

### Pilot Results with Global Open Data


Country	Population Data Set	Total Rural Population	Rural Population within 2km of Roads	% of Rural Population within 2 km of Roads
Nigeria 1	GRID3	86,073,240	80,056,026	93.0
	HRSL	91,823,234	84,610,846	92.1
	GHS Pop	49,553,503	45,167,977	91.1
Nigeria 2	GRID3 HRSL GHS Pop	86,073,240 91,823,234 49,553,502	55,136,377 59,071,979 33,317,594	64.1 64.3 67.2
Colombia	HRSL GHS Pop	21,304,160 7,589,940	18,396,019 6,894,459	86.3 90.8
Spain	HRSL GHS Pop	14,892,604 11,072,657	14,676,806 10,960,981	98.6 99.0

RAI results for Nigeria (1, with "unclassified" roads; 2, without "unclassified" roads), Colombia, and Spain using different gridded population inputs.

Source: Rural Access to Roads StoryMap on SDGs Today https://arcg.is/04PaHv

NASA's Applied Remote Sensing Training Program


#### Global Open Data RAI Results: Nigeria




50-75 %

25-50 %

### National RAI Results: CIESIN Roads Definition






\* Akrotiri and Dhekelia, Kosovo, Aland, and Northern Cyprus are excluded from this map because they do not have official UN population values for verifying their results.


\*\* Countries with a population percent error greater than 50 % when compared to official UN population values are excluded from this map.

0 - 20 20 - 40 40 - 60 60 - 80 80 - 100 RAI Not Computed

48

### National RAI Results: World Bank Roads Definition






\* Akrotiri and Dhekelia, Kosovo, Aland, and Northern Cyprus are excluded from this map because they do not have official UN population values for verifying their results.

\*\* Countries with a population percent error greater than 50 % when compared to official UN population values are excluded from this map.

0 - 20 20 - 40 40 - 60 60 - 80 80 - 100 RAI Not Computed

## Africa Subnational (L2) RAI Results: CIESIN Roads Definition




N = 8,436

#### 📕 0 - 20 📕 20 - 40 📕 40 - 60 🔜 60 - 80 🦳 80 - 100 📰 RAI Not Computed



## Africa Subnational (L2) RAI Results: World Bank Roads Definition



N = 9,364

#### 📕 0 - 20 📕 20 - 40 📕 40 - 60 🥅 60 - 80 🦳 80 - 100 📰 RAI Not Computed



# **Preliminary Global Average RAI Results**

#### World Bank Roads Definition

|--|

	GHS	HRSL	WP
AVERA			
GE	83.4	82.9	76.4
STDEV	18.2	16.7	21.1

	HRSL	
92.4	93.8	86.7
12.3	9.1	15.5
		92.4 93.8

CIESIN'22npre inclusive select2200f "all-seasen" roads resul222global apperage R2221 that are consistently ~10 percentage points greater than the averages computed with the World Bank definition.

- For both definitions, GHS & HRSL averages are generally consistent with each other and ~8 points higher than the WP results.
- This is consistent with WP often predicting lower RAIs for a given unit, likely due to the the manner in which the model allocates population to large rural areas.





Gridded Population Estimates	<ul> <li>Several data sets to choose from, each with their own bias</li> <li>Varying frequency of updates</li> </ul>
GADM National Boundaries	<ul> <li>Resolution varies by country</li> <li>Handling of disputed territories</li> <li>Infrequent/unpredictable updates</li> </ul>
S-MOD Urban/Rural Areas	<ul> <li>Potential misalignment with national urban/rural delineation</li> </ul>
OSM Roads Accuracy ultimately dep NASA's Applied Remote Sensing Training Proc	<ul> <li>Variable coverage, especially in rural areas         <ul> <li>Limited road condition information</li> <li>Imited road condition</li> <li>Imited road</li></ul></li></ul>

### Let's Review...

- SDG indicator 9.1.1, the **Rural Access Index** (RAI), measures **rural access to all-season roads**.
- RAI can be effectively calculated by combining geospatial data for **administrative units**, **rural areas**, **roads**, and **population**.
- While there are **advantages** to using global datasets, such as standard inputs, automation, and temporal comparisons, there are also **limitations**.
- The choice of **POPGRID** dataset and **OSM road selections** significantly impact RAI results calculated using this method.
- This is a simple and effective method for monitoring RAI globally.



## **References:**

Roberts, Peter; KC, Shyam; Rastogi, Cordula. 2006. Rural Access Index : A Key Development Indicator. Transport paper series; no. TP-10. World Bank, Washington, DC. © World Bank. https://openknowledge.worldbank.org/handle/10986/17414 License: CC BY 3.0 IGO.

Transport & ICT. 2016. Measuring Rural Access: Using New Technologies. Washington DC: World Bank, License: Creative Commons Attribution CC BY 3.0 <a href="http://documents.worldbank.org/curated/en/367391472117815229/Measuring-rural-access-using-new-technologies">http://documents.worldbank.org/curated/en/367391472117815229/Measuring-rural-access-using-new-technologies</a>

World - Measuring Rural Access : Update 2017/18 (English). Washington, D.C. : World Bank Group. http://documents.worldbank.org/curated/en/543621569435525309/World-Measuring-Rural-Access-Update-2017-18

Schiavina, Marcello; Freire, Sergio; MacManus, Kytt (2019): GHS population grid multitemporal (1975, 1990, 2000, 2015) R2019A. European Commission, Joint Research Centre (JRC) DOI: <u>10.2905/42E8BE89-54FF-464E-BE7B-BF9E64DA5218</u> PID: <u>http://data.europa.eu/89h/0c6b9751-a71f-4062-830b-43c9f432370f</u>

Pesaresi, Martino; Florczyk, Aneta; Schiavina, Marcello; Melchiorri, Michele; Maffenini, Luca (2019): GHS settlement grid, updated and refined REGIO model 2014 in application to GHS-BUILT R2018A and GHS-POP R2019A, multitemporal (1975-1990-2000-2015), R2019A. European Commission, Joint Research Centre (JRC) DOI: 10.2905/42E8BE89-54FF-464E-BE7B-BF9E64DA5218 PID: http://data.europa.eu/89h/42e8be89-54ff-464e-be7b-bf9e64da5218

Worldpop. "Open Spatial Demographic Data and Research." WorldPop. Accessed February 03, 2021. https://www.worldpop.org/.

"Methodology: High Resolution Population Density Maps Demographic Estimates." Facebook Data for Good. February 02, 2020. Accessed February 04, 2021. https://dataforgood.fb.com/docs/methodology-high-resolution-population-density-maps-demographic-estimates/.

"SDG Indicators - SDG Indicators." United Nations. Accessed February 04, 2021. https://unstats.un.org/sdgs/metadata/?Text=&Goal=9&Target=9.1.

"GADM Maps and Data." GADM. Accessed February 04, 2021. https://gadm.org/.

"Global Openstreetmap Data Extracts." Osmdata.xyz. Accessed February 04, 2021. https://download.osmdata.xyz/.


"Our Download Server." GEOFABRIK // Downloads. Accessed February 15, 2021. https://www.geofabrik.de/data/download.html.

"High Resolution Settlement Layer." Center for International Earth Science Information Network. Accessed March 08, 2021. https://ciesin.columbia.edu/data/hrsl/.

Worldpop. "WorldPop Methods." WorldPop. Accessed March 08, 2021. https://www.worldpop.org/methods.

"Global Population Grids: Summary Characteristics." Home. Accessed March 08, 2021. https://www.popgrid.org/data-docs-table1.







# 2021 ARSET-Applied Sciences Disasters Program

Population Grid Use Cases: Modeling Building Exposure data for DRR Charles K. Huyck, Co-Founder ImageCat

March 30th and April 6th , 2021

# Funding and Background

- Research funded by UK Space Agency International Partnership Programme: Project METEOR
- And through the NASA Disasters Program: Open Critical Infrastructure Exposure for Disaster Forecasting, Mitigation, and Response
- And Human Planet funding from NASA









# What you should learn in this lesson...

- What is exposure data and how is it used in the process of estimating losses from natural disasters?
- The basic process of developing exposure data
- The value of EO data
- Understanding uncertainty and the appropriate use of results





# Loss Estimation and CAT (Catastrophe) Modeling for DRR

27

- **Before an Event -** What might happen if ...?
- During an Event Where is this going? What should we do?
- After an Event What just happened? Should we ask for international help? Where is the most damage? Where are people without food and shelter?
- On Average Where should we build stronger, higher, or farther away? Where should we retrofit, acquire property, or replace facilities? What should be insured? (Insurers- how much should that cost?)






# What is Building Exposure for DRR?






### Risk - Confluence of Hazard, Vulnerability, and Exposure





NASA's Applied Remote Sensing Training Program


# What is Building Exposure for DRR?



62


NASA's Applied Remote Sensing Training Program


# Number of buildings, where, by type and cost, for the purpose of assessing vulnerability and hazard proximity:





## How is exposure data developed?







NASA's Applied Remote Sensing Training Program


# **Estimating Building Attributes**

- Literature review of predominant building construction types
- Interpretation of EO
- Expert opinion
- Virtual reconnaissance
- Site surveys/stratified sampling
- Housing census



# Estimating Number and Size of Buildings by Development Pattern

- People per household and households per building
- OSM (Open Street Map) building data footprint
- Microcensus or government reports
- Distribution of building area by story height
- Population density, urbanity, or EO metrics sometimes correlate





## **Estimate Replacement Value**


- Building value per meter by building type or occupancy
- Building construction manuals
- Expert opinion
- GDP/median income
- Scale by building durability





# **Building Vulnerability Attributes**


- Era of development
- Number of stories
- First floor elevation
- Construction materials
- Lateral force resisting system
- Retrofitting
- Nail density
- Distance between buildings






# What is the role of EO in developing exposure data?

- Global population datasets
- Global urban/rural or urban intensity datasets
- Segmentation of development patterns
- Building footprint extraction
- Average building size
- Al, ground-based sensors




# **Building Footprint Extraction**





## **Building Footprint Extraction**





#### Residential



#### High Density Residential






# **Extracting Development Patterns**


Remote Sensing Example: Typical Midrise Residential in China


- Large residential apartments up to eight stories
- Typically constructed of unreinforced brick walls with concrete floor and roof diaphragms
- Observed RC frame with URM infill in newer buildings as well as typical Soviet Bloc construction












#### **Extracting Development Patterns**

## 27

More Accurate




**Higher Resolution** 




NASA's Applied Remote Sensing Training Program

#### Merging Structural Characteristics with Development Patterns






#### How good is this stuff?



#### "Law of Large Numbers"










#### (Whoops!)






30


NASA's Applied Remote Sensing Training Program

#### Levels of Exposure Data






#### Levels of Exposure Data



#### Levels of Exposure Data








### Key, Often Ignored Factors

- Persons per household
- Living area per household
- Rebuilding cost
- Exchange rates

#### Focus:

- Vulnerability
- Geographic resolution


## Global Population Datasets - Variance might not be critical for regional disasters.



Credit: Cascade Tuholske, PhD Postdoctoral Research Scientist Center for International Earth Science Information Network (CIESIN) Earth Institute | Columbia University www.tuholske.com



#### **Extracting Development Patterns**





#### Important to Address Expectations

Cannot typically expect accurate number of buildings at the cell level

- Can expect more accuracy than in the original base data sets
- Cannot expect to always capture small, unmapped rural areas
- Challenges in remote sensing that will impact results (low lights, cloud cover, tree canopies, etc.)
- Can not repurpose the data for civic purposes such as address-specific information for tax purposes











# How much does it cost to build this hut?



Why does it matter?

YES

YES Is it cost effective to retrofit certain types of buildings regionally? Where should we focus retrofitting efforts? Are building codes cost effective, and where? What might happen after...

A hundred-year flood

A large earthquake

A volcano

There has just been a large earthquake... What are the likely impacts?

Where is likely to have been affected the most?

How should we deploy resources?

Is it cost effective to retrofit <u>this</u> building? Which buildings fell down? Which homes are flooded? Exactly how many buildings fell down?









#### **Thank You!**



NASA's Applied Remote Sensing Training Program